Reg No.: Name:

# APJ ABDUL KALAM TECHNOLOGICAL UNIVERSITY

SECOND SEMESTER B.TECH DEGREE EXAMINATION, JULY 2018

**Course Code: MA102** 

**Course Name: DIFFERENTIAL EQUATIONS** 

**Duration: 3 Hours** 

Max. Marks: 100

PART A

Answer all questions, each carries 3 marks

Consider the initial value problem  $y'' - x^3y' + 6xy = \sin x$ , y(0) = 3, y'(0) = -1.

Can this problem have unique solution in an interval containing zero? Explain.

Find any three independent solutions of the differential equation y''' - y' = 0. (3)

- Find the particular solution of the differential equation  $y'' 6y' + 9y = e^{3x}$ . (3)
- Using a suitable transformation, convert the differential equation  $(2x-3)^2 y'' (2x-3)y' + 2y = (2x-3)^2 \text{ into a linear differential equation with constant coefficients.}$
- State the conditions for which a function f(x) can be represented as a Fourier series. (3)
- Discuss the convergence of a Fourier series of a periodic function f(x) of period  $2\pi$ .
- Find the partial differential equation representing the family of spheres whose (3) centers lies on z-axis.
- Find the particular solution of  $(D^2 2DD' + 2D'^2)z = \sin(x y)$  (3)
- 9 Write any three assumptions involved in the derivation of one dimensional wave (3) equation.
- A string of length l fastened at both ends. The midpoint of the string is taken to a height h and then released from rest in that position. Write the boundary conditions and initial conditions of the string to find the displacement function y(x,t) satisfying the one dimensional wave equation.
- Write the fundamental postulates used in the derivation of one dimensional heat (3) equation.
- Define steady state condition in one dimensional heat equation  $\frac{\partial u}{\partial t} = \alpha^2 \frac{\partial^2 u}{\partial x^2}$ . (3)

### **PART B**

# Answer six questions, one full question from each module

# **Module 1**

13 a) Discuss the existence and uniqueness of solution of the initial value problem (6)

$$\frac{dy}{dx} = \frac{y}{\sqrt{x}}, y(1) = 3.$$

b) Prove that  $y_1(x) = e^x$  and  $y_2(x) = e^{4x}$  form a fundamental system(basis) for the (5)

differential equation y'' - 5y' + 4y = 0. Can  $5e^x - 2e^{4x}$  be a solution (do not use verification method) of the differential equation? Explain.

Discuss the existence and uniqueness of solution of the initial value problem 14 (6) $\frac{dy}{dx} = x^2 + y^2$ , y(0) = 1 in the rectangle  $|x| \le 1, |y - 1| \le 1$ .

b) If  $y_1(x)=x$  is a solution of  $x^2y''+2xy'-2y=0$ , find the general solution. (5)

# **Module II**

By the method of variation of parameters, solve  $y'' + y = x \sin x$ . 15 (6)

b) Solve  $y'' + 5y' + 6y = e^{-2x} \sin 2x$ . (5)

### OR

16 a) Solve  $x^2y'' + xy' - 9y = \log x$ . (6)

b) Solve  $v'' - 2v' + 5v = x^2$ . (5)

### **Module III**

17 Find the Fourier cosine series representation of  $f(x)=x, 0 \le x \le \pi$ . Also find the (11)Fourier series representation f(x) if f(x) is periodic function with period  $\pi$ .

Find the Fourier series of the periodic function f(x) of period 4, where 18  $f(x) = \begin{cases} 2, & -2 < x \le 0 \\ x, & 0 < x < 2 \end{cases}$  and deduce that

(i)  $1 + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \dots = \frac{\pi^2}{8}$  and (ii)  $1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \dots = \frac{\pi}{4}$ 

Find the particular solution of  $\frac{\partial^2 z}{\partial x^2} + 3 \frac{\partial^2 z}{\partial x \partial y} + 2 \frac{\partial^2 z}{\partial y^2} = y^2$ . 19 (5)

Find the general solution of  $(y^2 + z^2)p - xyq = -xz$ . (6)

20 a) Solve  $(D^2 + 3DD' + 2D'^2)z = (2x + y)^7$ . (5)

(6)Solve  $4\frac{\partial^2 z}{\partial x^2} - 4\frac{\partial^2 z}{\partial x \partial y} + \frac{\partial^2 z}{\partial y^2} = 16 \log(x + 2y)$ .

## Module V

21 (5) Using method of separation of variables, solve  $\frac{\partial u}{\partial x} = 2 \frac{\partial u}{\partial t} - u$ ,  $u(x,0) = 5e^{-3x}$ .

(5) A tightly stretched string of length l fastened at both ends is initially in a position given by y = kx, 0 < x < 1. If it is released from rest from this position, find the displacement y(x,t) at any time t and any distance x from the end x = 0.

22 A string is stretched and fastened in two points 50 cm apart. Motion is started by displacing the string into the form of the curve y = x(50 - x) and also by imparting a constant velocity V to every point of the string in the position at time t = 0. Determine the displacement function y(x, t).

# **Module VI**

A rod of length 50 cm has its ends A and B kept at  $20^{\circ}$ C and  $70^{\circ}$ Crespectively (10) until steady state temperature prevail. The temperature at each end is thensuddenly reduced to zero temperature and kept so. Find the resulting temperature function u(x,t) taking x=0 at A.

### OR

A bar 10 cm long with insulated sides has its ends A and B maintained t 50°C (10) and 100°C respectively until steady state conditions prevail. The temperature at A is suddenly raised to 90°C and at the same time that at B is lowered to 60°C. Find the temperature distribution in the bar at time t.

\*\*\*\*

