	-

# B1A005

Pages: 3

Reg. No.\_\_\_\_\_\_ Name:\_\_\_\_\_

## APJ ABDUL KALAM TECHNOLOGICAL UNIVERSITY

FIRST SEMESTER B.TECH DEGREE EXAMINATION, JUNE/JULY 2017

Course Code: MA 101
Course Name: CALCULUS

## (For 2015 Admission and 2016 Admission)

Max. Marks:100 Duration: 3 hours

## PART A

## Answer all questions. Each question carries 5 marks.

- 1. (a) Find the interval of convergence and radius of convergence of the infinite series  $\sum_{n=0}^{\infty} n! \, x^n$  (2)
  - (b) Determine whether the series  $\sum_{n=1}^{\infty} \frac{1}{5n-1}$  converges or not (3)
- 2. (a) Find the slope of the surface  $z = \sqrt{3x + 2y}$  in the y-direction at the point (4,2)
  - (b) Find the derivative of  $w = x^2 + y^2$  with respect to t along the path  $x = at^2$ , y = 2at(3)
- 3. (a) Find the directional derivative of  $f(x, y) = xe^y$  at (1,1) in the direction of the vector i j (2)
  - (b) If  $\vec{F}(t)$  has a constant direction, then prove that  $\vec{F} \times \frac{d\vec{F}}{dt} = 0$  (3)
- 4. (a) Evaluate  $\int_0^1 \int_0^1 \frac{1}{\sqrt{(1-x^2)(1-y^2)}} dxdy$  (2)
  - (b) Evaluate  $\iint_R \frac{\sin x}{x} dxdy$  where R is the triangaular region bounded by the x-axis, y = x and x = 1.
- 5. (a) Show that  $\int_A^B (2xy + z^3) dx + x^2 dy + 3xz^2 dz$  is independent of the path joining the points A and B. (2)
  - (b) If  $\vec{r} = xi + yj + zk$  and  $r = |\vec{r}|$ , then prove that  $\nabla^2 r^n = n(n+1)r^{n-2}$  (3)
- 6. (a) Using line integral evaluate the area enclosed by the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  (2)
  - (b)Evaluate  $\int_C (e^x dx + 2y dy dz)$  where C is the curve  $x^2 + y^2 = 4$ , z = 2. (3)

## PART B

### Answer any two questions each Module I to IV

#### Module I

- 7. Determine whether the series converge or diverge  $\sum_{n=1}^{\infty} \frac{(2n)!}{(n!)^2}$  (5)
- 8. Check the absolute convergence or divergence of the series  $\sum_{n=1}^{\infty} (-1)^n \frac{(2n-1)!}{3^n}$  (5)

9. Find the Taylor series expansion of  $\log \cos x$  about the point  $\frac{\pi}{3}$  (5)

#### Module II

10. If 
$$u = log (x^3 + y^3 + z^3 - 3xyz)$$
, Show that  $\left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} + \frac{\partial}{\partial z}\right)^2 u = \frac{-9}{(x+y+z)^2}$  (5)

- 11. The length, width and height of a rectangular box are measured with an error of atmost 5%. Use a total differential to estimate the maximum percentage error that results if these quantities are used to calculate the diagonal of the box. (5)
- 12. Locate all relative extrema and saddle points of  $f(x, y) = x^4 + y^4 2x^2 + 4xy 2y^2$  (5)

#### Module III

- 13. Find the angle between the surfaces  $x^2 + y^2 + z^2 = 9$  and  $z = x^2 + y^2 3$ at the point (2, -1, 2)
- 14. Let  $\vec{r} = xi + yj + zk$  and  $r = |\vec{r}|$ , then prove that  $\nabla f(r) = \frac{f'(r)}{r}\vec{r}$ . (5)
- 15. Find an equation of the tangent plane to the ellipsoid  $2x^2 + 3y^2 + z^2 = 9$ at the point (2,1,1) and determine the acute angle that this plane makes with the XY plane. (5)

### **Module IV**

- 16. Change the order of integration and hence evaluate  $\int_0^1 \int_{x^2}^{2-x} xy \, dy \, dx$  (5)
- 17. Evaluate  $\int_0^2 \int_0^{\sqrt{(4-x^2)}} y(x^2 + y^2) dx dy$  using polar co-ordinates (5)
- 18. Find the volume of the paraboloid of revolution  $x^2 + y^2 = 4z$  cut off by the plane z = 4 (5)

## Module V

### Answer any 3 questions.

- 19. Evaluate the line integral  $\int_C (xy + z^3) ds$  from (1,0,0) to (-1,0, $\pi$ ) along the helix C that is represented by the parametric equations  $x = \cos t$ ,  $y = \sin t$ , z = t (5)
- 20. Evaluate the line integral  $\int_C (y-x) dx + x^2 y dy$  along the curve C,  $y^2 = x^3$  from (1, -1) to (1,1) (5)
- 21. Find the work done by the force field  $\vec{F} = (x + y)i + xyj z^2k$  along the line segment from (0,0,0) to (1,3,1) and then to (2,-1,5).
- 22. Show that  $\vec{F} = (2xy + z^3)i + x^2j + 3xz^2k$  is a conservative vector field. Also find its scalar potential. (5)
- 23. Find the values of constants a, b, c so that  $\vec{F} = (axy + bz^3)i + (3x^2 cz)j + (3xz^2 y)k$  may be irrotational. For these values of a, b, c find the scalar potential of  $\vec{F}$  (5)

#### Module VI

# Answer any 3 questions.

- 24. Verify Green's theorem for  $\int_C (xy + y^2)dx + x^2 dy$  where C is bounded by y = x and  $y = x^2$  (5)
- 25. Apply Green's theorem to evaluate  $\int_C (2x^2 y^2)dx + (x^2 + y^2)dy$  where C is the boundary of the area enclosed by the x-axis and the upper half of the circle  $x^2 + y^2 = a^2$ (5)
- 26. Apply Stokes theorem to evaluate  $\int_C (x+y)dx + (2x-y)dy + (y+z)dz$  where C is the boundary of the triangle with vertices (0,0,0), (2,0,0) and (0,3,0) (5)
- 27. Use Divergence theorem to evaluate  $\iint_S \vec{F} \cdot \vec{n} dS$  where  $\vec{F} = xi + zj + yzk$  and S is the surface of the cube bounded by x = 0, x = 1, y = 0, y = 1, z = 0 and z = 1. Also verify this result by computing the surface integral over S (5)
- 28. State Divergence theorem. Also evaluate  $\iint_S \vec{F} \cdot \vec{n} dS$  where  $\vec{F} = axi + byj + czk$  and S is the surface of the sphere  $x^2 + y^2 + z^2 = 1$  (5)

\*\*\*